

Research Article

Life cycle study of maize stem borer (*Chilo partellus* Swinhoe) under laboratory condition at National Maize Research Program, Rampur, Chitwan, Nepal

¹Saraswati Neupane * and ¹Subash Subedi,

¹Nepal Agricultural Research Council, National Maize Research Program Rampur, Chitwan, Nepal

*Correspondence: sarusanu2017@gmail.com; ORCID: <https://orcid.org/0000-0003-3033-5840>

Received: August 18; Accepted: October 5; Published: October 25, 2019.

© Copyright: Neupane and Subedi (2019).



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

ABSTRACT

Maize stem borer (*Chilo partellus* Swinhoe) is one of the major threatening global pests of maize and considered as the national top priority entomological research problem in Nepal. The Life cycle of maize stem borer was studied under laboratory condition at National Maize Research Program (NMRP), Rampur, Chitwan, Nepal during 2018. Development of stem borer undergoes following stages like egg, larvae, pupa and adult. Eggs and different instars of maize stem borer larvae were collected from maize fields were put with host materials (maize leaf and stem) to become different instars of larva, pupae and finally turned to adults. Eggs were harvested from adults and kept on blotting paper which was kept inside petriplates and reared for adults. Their life span in each stage (egg, larva and pupa) and the fecundity of adults recorded. Daily room temperature and relative humidity (RH) in laboratory conditions were recorded. The Egg incubation period ranged from 4-7 days and hatched generally in the early morning (6-8 AM). The complete larvae period ranged from 29 to 36 days while pupal period was ranged from 7 to 12 days. The average male pupal length was found 10.6 mm and female was 13.13 mm long. The fecundity of *C. partellus* Swinhoe was recorded 150-160 eggs per female. The Oviposition period was 4-5 days and adult male survived for 4-7 days while female for 5-9 days. The average life cycle of *C. partellus* completed in (44-48) days during summer whereas (60-64) days during winter at average room temperature of (26-27° C) and RH of (70-80%). These results have important implications to know the survival and development of pest including effective pest management strategy.

Keywords: *Chilo partellus* Swinhoe, life cycle, maize

Correct citation: Neupane, S., & Subedi, S. (2019). Life cycle study of maize stem borer (*Chilo partellus* Swinhoe) under laboratory condition at National Maize Research Program, Rampur, Chitwan, Nepal. *Journal of Agriculture and Natural Resources*, 2(1), 338-346.

DOI: <https://doi.org/10.3126/janr.v2i1.26099>

INTRODUCTION

Insect pests are one of the major biotic constraints that limit the productivity of maize in Nepal. More than 130 insects have been reported to cause damage to maize and about half a dozen insects are economically importance (Atwal, 1976). The major insect pests that attack maize crop include maize aphids, cutworms, stem/shoot fly, root worm and stem borer (Thakur *et al.*, 2018). Among them, maize stem borer, *Chilo partellus* Swinhoe (Pyralidae: Lepidoptera) has been implicated as a major production constraint throughout the country (Neupane, 1986; Sharma & Gautam, 2010; Thakur *et al.*, 2018). It attacks maize plants from seeds sown in field to maturity and feed on all parts of the plants (Gyawali, 1978; Shivakoti & KC, 1981; Neupane *et al.*, 1983). Newly hatched larvae feed on the leaves making pinholes and leaf windowing. They bore down inside the plant whorl and feed. While feeding in the plant whorl, they kill the central shoot, which later on dries up causing dead heart resulting total loss of the crop (Harris, 1990). In general, the borer caused minor damage, pin holes on leaves. In severe cases, dead hearts are formed, the condition in which the plants do not bear any ear at all. At our context, till now, yield loss caused by maize stem borer yet to be estimated in combined with the relationship of stem tunneling as well as leaf damage to monetary value. However, some researchers have reported that borer can culminate yield loss of 20-87% (Mathur, 1994; Chatterji *et al.*, 1969). The figure of on-farm of yield loss caused by borer is considerable. It is very difficult to manage stem borer by chemical insecticides alone due to their ability to develop quick resistance against insecticides. The biological study of any pest is always important to explore the survival and development knowhow of the pest and ultimately formulate the effective and sustainable pest management strategy. The time and mode of pesticide application as well as pest behaviour towards host during infection process warrant the information of pest biology otherwise spraying of chemical during egg stage, late larval and pupal stage is ineffective. Scanty information is available on biological aspect of this pest in Nepal. Considering and realizing the significance of these aspects, to know the detail life cycle pattern of maize stem borer, the present studies were under taken at the laboratory of National Maize Research Program, Rampur, Chitwan, Nepal.

MATERIALS AND METHODS

The Life cycle of maize stem borer was studied under laboratory condition at National Maize Research Program (NMRP), Rampur, Chitwan, Nepal during 2018. Development of stem borer undergoes following stages like egg, larvae, pupa and adult. Daily room temperature and relative humidity (RH) in laboratory conditions were recorded.

Maintenance of insect culture

Eggs and different instars of maize stem borer were collected from maize field at National Maize Research Program, Rampur, Chitwan. The collected larvae were transferred to plastic boxes containing cut stems of maize. The boxes were cleaned daily by removing the excreta and partially eaten stems. The larvae were supplied with sufficient quantity of fresh tender stem pieces as food. The larvae were reared till they attain pupal stage. Pupae obtained were kept in separate plastic jar for adult emergence (Arunkumara *et al.*, 2018). The culture then obtained was used for the further studies on biology conducted in growth chamber at 27 °C temperature and RH of 75 per cent.

Incubation period

For this five pairs of freshly emerged male and female adults were released into oviposition jar (25 x 10 cm) which was prepared by covering the inner wall of jar with folded paper, fresh maize plants. Ten per cent honey solution was provided as a food to the moths by soaking cotton plug in the solution. The open end of oviposition jar was covered with muslin cloth. Folded paper and leaves served as an ovipositional site and observations were recorded every day for oviposition. Eggs laid on the folded paper by the female moth were transferred with the help of a camel hair brush and needle to a petri-dish with moistened cotton sponge. The petridish was covered and observed for hatching of the egg. Time taken for hatching was recorded by making observation at 24 hours interval.

Larval period

The period from hatching of the eggs till pupation was recorded as larval period. Larvae reared on fresh maize leaf and stem in laboratory condition. These larvae were observed for the morphometric characters (Arunkumara *et al.*, 2018).

Pupal period

Pupae obtained after completion of the larval cycle were collected in plastic container and kept for emergence. The time taken from the pupation till emergence of the adult recorded as a pupal period. These pupae were observed for the morphometric characters (Arunkumara *et al.*, 2018).

Adult longevity

Adult emerged from pupa was observed for their colour, shape and size. The male and female moth was measured across their expanded wing with the help of standard scale. Male and female were differentiated on the basis of morphological differences. The male and female longevity were calculated separately from date of emergence of adult to the death of adult.

Oviposition period

The period from first egg laid till the last egg laid was recorded as oviposition period.

Fecundity

Individual pairs of freshly emerged moths were kept in cylindrical boxes provided with folded paper and 10% honey solution. Honey and oviposited paper were changed daily till the death of female of the pair. Eggs were counted on oviposited paper and leaves on daily basis and finally total fecundity was calculated.

Total life cycle

The duration of entire life span of insect was considered as total life cycle.

Data analysis

All data were analyzed statistically using Microsoft Excel (2007) computer package program. The average mean, standard deviation and standard error was calculated.

RESULTS AND DISCUSSION

Incubation period of egg

The eggs were flat oval and creamy white in color. An eggmass consisted of 20-35 eggs, which were overlap and scale like in appearance (Figure 1). The higher incubation period (6-7 days) was observed during September to February while lower period (4-5 days) during May to August (Table 1). At initial stages egg get appeared from pearly white in color and then finally turns to darker in color. Egg hatched in 4-7 days generally in the early morning (6 am-8 am) (Table 4). This finding is in line with the Krishna *et al* (2018) who reported that eggs of stem borer get hatched within 5-6 days during early morning were flat and creamy white color and finally turns to darker color which represent the head of the larvae.

Larvae

The larvae was found with black or dark brown colored head, purplish pink on dorsal side and white on ventral side (Figure 2). The average mean value of length, breadth and weight of the first and fifth instar larvae observed from 2.05 ± 0.11 to 23.38 ± 0.55 mm; 0.34 ± 0.03 to 3.22 ± 0.17 mm and 1.04 ± 0.06 to 327.5 ± 43.3 mg respectively (Table 3). The higher larval period (35-36 days) was recorded during the month January-February while lower (29-30 days) in May-June (Table 1). The total larval period was ranged from 29-36 days with the mean value of 32.28 ± 0.17 (Table 4). In the larval period mean value of instars I, II, III, IV and V observed from 4.90 ± 0.28 , 6.50 ± 0.17 , 8.40 ± 0.16 , 9.30 ± 0.15 and 10.40 ± 0.16 respectively (Table 4). The present findings with respect to larval duration are in agreement with that of Deshpande (1978) and Marulasiddesha (1999). The morphometric measurements of the pest are also in line with the findings of Krishna *et al* (2018).

Pupae

The pupae were pale brown with cylindrical shape (Figure 3). The higher pupal period (11-12 days) was recorded during the month November-February while lower period (7-8 days) in May-June (Table 2). The male pupae was slightly smaller than that of female pupae. The average mean length, breadth and weight of the male pupae observed from 10.66 ± 0.23 mm, 2.23 ± 0.07 mm and 29.15 ± 1.28 mg respectively (Table 3). In case of female pupae, the average mean length, breadth and weight were 13.13 ± 0.15 mm, 2.66 ± 0.02 mm and 49.15 ± 0.74 mg respectively (Table 3). The male pupal period ranged from 7-9 days with the mean value of 7.70 ± 0.26 days while 10-12 days with average mean value of 10.90 ± 0.23 days for female pupae (Table 4). The present findings are in close agreement with Deshpande (1978), Krishna *et al* (2018) and Arunkumara *et al* (2018) who reported that pupal period lasted for 6 to 11 days.

Adult

The adults were medium sized with yellowish brown forewings having rows of black spots all along anterior margin. Male moth was dark brown in colour with pale brown forewings and pale yellow hindwing while female hind wing was much lighter than that of male (Figure 4). The higher adult period (8-9 days) was recorded during the month January-February while lower (4-5 days) in May-June (Table 2). The average mean length, breadth and weight of the male adult observed from 13.53 ± 0.22 mm, 23.55 ± 0.22 mm and 49.12 ± 0.29 mg respectively (Table 3). In case of female pupae, the average mean length, breadth and weight were 18.55 ± 0.23 mm, 29.84 ± 0.32 mm and 57.62 ± 0.51 mg respectively (Table 3). The male adult period ranged from 4-7 days with the mean value of 5.40 ± 0.34 days while 5-9 days with average mean value of 7.0 ± 0.39 days for female adult (Table 4). The finding from

this study is well supported by Berger (1992) , Bleszynski (1970) and Harris (1990) who reported that adults were relatively small moths with wing lengths ranging from 7 to 17 mm. Females are generally larger than males and both sexes rest with the wings folded over the abdomen. The forewings are generally light yellow-brown with some darker scale patterns forming longitudinal striations which are usually darker at the wing margins (Arunkumara *et al.*, 2018).

Oviposition period

The oviposition period is the period from first egg laid till the last egg laid and it ranged from 4-5 days with average mean value of 4.2 ± 0.13 days (Table 4). These results are in the conformity with earlier records of Chavan (2006) who reported that oviposition period of 3 to 5 with an average of (4.18 ± 0.64) days. Similar, another report also reported the oviposition period to ranges from 3.00 to 5.00 days with an average of 4.2 ± 0.63 days. Thus, the present finding of oviposition period was more or less in conformity with past report reported by Siddalingapa *et al.* (2010).

Fecundity

The fecundity of *C. partellus* Swinhoe was recorded 150-160 eggs per female with the mean value of 157 ± 0.52 eggs per female (Table 4). The findings of this study were in support with the findings of Songa *et al* (2000) and Berner *et al* (1993) who reported that the fecundity of *C. partellus* was approximately 150 eggs per female.

Total life cycle

The average life cycle of *C. partellus* from egg to adult (Figure 5) completed in (44-48) days during peak summer period (May-June) whereas (60-64) days during peak winter season i.e. in January- February (Table 2). The total life cycle period ranged from 44-64 days with the mean value of 53.82 ± 0.49 days (Table 4). These findings are in agreement with previous records of Marulasiddesha (1999) reported that life cycle of stem borer as 30 to 65 days and also reported the life cycle as 30 to 69 days with an average of 49.50 ± 13.50 days by Siddalingapa *et al.* (2010) .

Table 1: Egg incubation and larval period of *Chilo partellus* (Swinhoe) at laboratory of NMRRP, Rampur under natural feeding condition in 2018-2019

S. N.	Month (2018-2019)	Incubation period of egg (days)		Larval period (days)	
		Range	Mean \pm SE	Range	Mean \pm SE
1	July-August	4-5	4.5 ± 0.16	30-31	30.2 ± 0.13
2	September-October	6-7	6.7 ± 0.15	33-34	33.5 ± 0.17
3	November-December	6-7	6.5 ± 0.17	34-36	34.7 ± 0.26
4	January-February	6-7	6.4 ± 0.16	35-36	35.4 ± 0.16
5	March- April	5-6	5.8 ± 0.13	30-31	30.6 ± 0.16
6	May-June	4-5	4.7 ± 0.15	29-30	29.3 ± 0.15

Note: † mean of 10 observations, SE- Standard error




Table 2: Pupal, adult and total life cycle period of *Chilo partellus* (Swinhoe) at laboratory of NMRP, Rampur under natural feeding condition in 2018-2019

S. N.	Month (2018-2019)	Pupal period (days)		Adult period (days)		Total life cycle (days)	
		Range	Mean \pm SE	Range	Mean \pm SE	Range	Mean \pm SE
1	July-August	8-9	8.4 ± 0.16	5-6	5.3 ± 0.15	47-51	48.7 ± 0.54
2	September-October	9-10	9.8 ± 0.13	5-7	6.0 ± 0.30	54-58	55.9 ± 0.48
3	November-December	11-12	11.3 ± 0.15	7-8	7.2 ± 0.13	59-63	60.7 ± 0.50
4	January-February	11-12	11.6 ± 0.16	8-9	8.4 ± 0.16	60-64	62.10 ± 0.50
5	March- April	8-9	8.8 ± 0.13	5-6	5.6 ± 0.16	48-52	49.70 ± 0.45
6	May-June	7-8	7.5 ± 0.17	4-5	4.7 ± 0.15	44-48	45.80 ± 0.47

Note: † mean of 10 observations, SE- Standard error

Table 3: Morphological measurements of *Chilo partellus* (Swinhoe) at laboratory of NMRP, Rampur under natural feeding condition in 2018-2019

S. N.	Growth stages	Length (mm)		Breadth (mm)		Weight (mg)	
		Range	Mean \pm SE	Range	Mean \pm SE	Range	Mean \pm SE
1	First instar larvae	1.5-2.5	2.05 ± 0.11	0.21-0.44	0.34 ± 0.03	0.8-1.4	1.04 ± 0.06
2	Second instar larvae	4.2-6.0	5.08 ± 0.21	0.73-0.85	0.8 ± 0.01	2.50-4.85	3.70 ± 0.32
3	Third instar larvae	8.2-15.5	12.48 ± 0.86	0.93-1.46	1.19 ± 0.06	10.6-30.45	22.49 ± 2.14
4	Fourth instar larvae	16.2-20.4	17.97 ± 0.44	1.75-2.54	2.18 ± 0.09	1.25-101.5	74.18 ± 5.30
5	Fifth instar larvae	20.6-25.2	23.38 ± 0.55	2.95-3.45	3.22 ± 0.17	110.5-490.6	327.5 ± 43.3
6	Male pupae	9.5-11.5	10.66 ± 0.23	1.95-2.52	2.23 ± 0.07	23.5-35.6	29.15 ± 1.28
7	Female pupae	12.5-13.8	13.13 ± 0.15	2.55-2.75	2.66 ± 0.02	45.5-52.5	49.15 ± 0.74
8	Male adult	12.5-14.5	13.53 ± 0.22	22.5-24.5	23.55 ± 0.22	47.5-50.4	49.12 ± 0.29
9	Female adult	17.5-18.5	18.55 ± 0.23	28.5-31.5	29.84 ± 0.32	55.5-60.5	57.62 ± 0.51


Note: † mean of 10 observations, SE- Standard error, mm- millimeter, mg-milligram

Table 4: Duration of different stages and fecundity of *Chilo partellus* (Swinhoe) at laboratory condition of NMRP, Rampur under natural feeding condition in 2018-2019

S.N.	Characteristics	Range	Mean \pm SE
1.	Egg incubation period (days)	4-7	5.75 ± 0.16
2.	Larval period (days)		
	2.1 First instar	4-6	4.90 ± 0.28
	2.2 Second instar	6-7	6.50 ± 0.17
	2.3 Third instar	8-9	8.40 ± 0.16
	2.4 Fourth instar	9-10	9.30 ± 0.15
	2.5 Fifth Instar	10-11	10.40 ± 0.16

3.	Total larval period (days)	29-36	32.28 ± 0.17
4.	Male pupal period (days)	7-9	7.70 ± 0.26
5.	Female pupal period (days)	10-12	10.90 ± 0.23
6.	Male adult (days)	4-7	5.4 ± 0.34
7.	Female adult (days)	5-9	7.0 ± 0.39
8.	Oviposition period (days)	4-5	4.2 ± 0.13
9.	Total life cycle period (days)	44-64	53.82 ± 0.49
10.	Fecundity (number)	155-160	157 ± 0.52

Note: † mean of 10 observations, SE- Standard error



CONCLUSION

The present findings from the study are important to understand the pest strategies during host infection process in order to formulate and design the sustainable pest management plan in the future. This study supplies ample information regarding the biology, survival, development and morphometric parameters of different stages of pest for developing pest resistant program.

ACKNOWLEDGEMENTS

The authors are grateful to all concerned scientists and technicians from the NARC stations supported during study period. Thanks are extended to the NMRP team for continuous support to carry out the experiment and facilities to prepare this manuscript. The funding for this study was received from NARC.

Author Contributions

S.N. designed and organized the experiment, analysed data and wrote the paper; S.S. helped during manuscript preparation.

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

REFERENCES

- Arunkumara, C.G., Bheemanna, M., Shaila, H.M., Teggelli, R., & Chavan, S. (2018). Biology of spotted stem borer *Chilo partellus* (Swinhoe) (Lepidoptera: Crambidae) on artificial diet. *Journal of Entomology and Zoology Studies*, 6(5), 685-688.
- Atwal, A.S. (1976). A Book of Agriculture Pests of India and South East Asia, 2nd Edition Kalyani Publisher, Delhi, India. p159.
- Berger, A. (1992). Larval movements of *Chilo partellus* (Lepidoptera: Pyralidae) within and between plants: timing, density responses and survival. *Bulletin of Entomological Research*, 82(4), 441-448
- Berner, D.K., Aigbokhan, E.I., & Ikie, F.O. (1993). Time of striga hermonthica infection in relation to parasite emergence and yield of sorghum and maize. *Phytopathology*, 83,13-63.
- Bleszynski, S. (1970). A revision of the world species of Chilo Zincken (Lepidoptera: Pyralidae). *Bulletin of the British Museum (Natural History)-Entomology*, 25:101-195.
- Chatterji, S.M., Young, W.R., Sharma, G.C., Sayi, I.V., Chall, B.S., Khare, B.P., Rathor, Y.S., Panwar, V.P.S., & Siddiqui, K.H. (1969). Estimate of loss in maize due to insect of maize with special reference to borer. *Indian Journal of Entomology*, 31, 109-115.
- Chavan, N.P. (2006). Biology, varietal screening against stem borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae) in sorghum (*Sorghum bicolor*) L. (Monech). M.sc. (Agri) thesis, Navsari Agricultural University, Navsari.
- Deshpande, V.P. (1978). Studies on the bionomics of sorghum stem borer, *C. partellus* (Swinhoe) and reaction of different sorghum varieties to it. M. Sc. (Agri.) Thesis, University of Agricultural Sciences Bangalore.
- Gyawali, B.K. (1978). Review of the entomological works on injurious maize pests in Nepal. The 5th National Maize Development Workshop. Naktajhij, Dhanusha. Pp: 166-188.
- Harris, K.M. (1990). Bioecology of Chilo species. *Insect Science and its Application*, 11(4-5), 467-477.

- Krishna, M.S.R., Peddakasim, D., Suneetha, P., Srideepthi, R., & Sahithya, U.L. (2018). Survival and development of maize stem borer *Chilo Partellus* (Swinhoe) Lepidoptera: Crambidae on artificial diet. *Acta Ecologica Sinica*, 38, 144–147.
- Marulasiddesha, K.N. (1999). Bio-ecology of stem borer, *Chilo partellus* (Swinhoe) and impact of its damage on juice quality of sweet sorghum. M.Sc. (Ag.) Thesis, University of Agricultural Sciences, Dharwad, Karnataka.
- Mathur, L.M.L. (1994). Advances in insect pest management in maize. In: G.S. Dhaliwal and Ramesh (eds.), Trends of agriculture insect pest management, common wealth publisher, New Delhi, India, pp. 113-160.
- Neupane, F.P. (1986). Temperature-developmental relationship of maize borer, *Chilo partellus* (Swinhoe) (Lepidoptera: Pyralidae), in Nepal. *Nepalese Journal of Agricultural Science*, 17, 23-35.
- Neupane, F.P., Chapman, R.K., & Coppel, H.C. (1983). Effect of date of planting of maize on the incidence of the maize borer, *Chilo partellus* (Swinhoe) in Nepal. *Journal of Institute of Agriculture & Animal Science*, 4, 77-79.
- Sharma, P.N. & Gautam, P. (2010). Assessment of yield loss in maize due to attack by the maize Borer, *Chilo partellus* (Swinhoe). *Nepal Journal of Science and technology*, 11, 25-30.
- Shivakoti, G.P. & K.C. Bhimsen. (1981). Screening maize germination against maize borer *Chilo partellus* Swinhoe. The ninth summer crops workshop held at National maize development program, Rampur, Chitwan, Nepal. Pp: 95-98.
- Siddalingappa, A., Thippeswamy, C., Venkatesh, H., & Shivsharanappa, Y. (2010). Biology of maize stem borer, *C. partellus* (Swinhoe) (Crambidae: Lepidoptera). *International Journal of Plant Protection*, 3(1), 91-93.
- Songa, J.M., Berginson, D., & Mugo, S. (2000). Mass rearing of the maize stem borers *Chilo partellus*, *Busseola fusca*, *Sesamia calamistis*, *Chilo orichalcociliellus* and *Eldana saccharina* KARI-Katamani. In proceedings of seventh Eastern and Southern Africa Regional maize conference 12-15 February 2001. pp. 120-124.
- Thakur, P., Bhandari, G., Shrestha, J., & Achhami, B. B. (2018). Effect of planting dates and varieties on infestation of maize stem borer *Chilo partellus* (Swinhoe). *International Journal of Applied Biology*, 2(1). 22-28